

# Depression

Major depressive disorder (MDD) is a serious mood disorder characterized by symptoms that interfere with an individual's feelings, thoughts, actions, and daily functioning. According to the World Health Organization, [depression](#) is the leading cause of disability worldwide, affecting more than 300 million individuals globally.

## Common types of depression

- **Bipolar disorder**, characterized by experiencing alternating periods of extremely high moods and extremely low moods
- **Persistent depressive disorder**, characterized by a depressed mood lasting over two years
- **Postpartum depression**, characterized by experiencing major depression during pregnancy and/or the postpartum period (directly following childbirth)
- **Psychotic depression**, characterized by severe depression with some form of psychosis (e.g., delusions, hallucinations)
- **Seasonal affective disorder (SAD)**, characterized by a cyclical onset of depression that is associated with less exposure to sunlight in the winter months

## Signs, symptoms, and complications

- Change in appetite and/or weight
- Decreased energy or [fatigue](#) (without significant exertion)
- Difficulty concentrating or remembering
- Disturbed [sleep](#)
- Feeling hopeless, guilty, worthless, or helpless
- Loss of pleasure in usual hobbies or activities
- Low or depressed [mood](#)
- Suicidal thoughts or suicide attempts

The severity, frequency, and duration of symptoms will vary depending on the individual. In order to be classified as MDD, some of the listed symptoms must be present almost every day for a minimum of two weeks.



# Causes and risk factors

- Being overweight or obese
- Certain chronic medical conditions (e.g., [dementia](#), Alzheimer's disease, epilepsy, Parkinson's disease, stroke, obesity, cancer, [arthritis](#))
- Certain medications (e.g., PPIs, antihypertensives, analgesics)
- Family history/genetic predisposition
- Gender: females have a greater risk of depression
- Major stress, trauma, or life events
- Socio-economic factors: poverty, housing difficulties, prejudice

# Preventing and addressing depression

While depression is typically addressed using a combination of medication, psychotherapy, and electroconvulsive therapy (in treatment-resistant cases), certain dietary and lifestyle interventions may also be beneficial.

## Diet

Research has shown that an overall dietary pattern focusing on whole foods while reducing processed or fast foods may reduce the risk of depression. The table below summarizes foods to avoid and enjoy in your diet.

Avoid	Enjoy
High-fat dairy products (e.g., butter, heavy cream)	Antioxidant-rich foods (e.g., dark chocolate, nuts, berries, leafy greens, herbs, spices)
Potatoes	Extra-virgin olive oil
Processed meat (e.g., hot dogs, cured meats, canned meat, sausages)	Fish (e.g., salmon, trout, mackerel)
Red meat (e.g., beef, pork, veal)	Fruit
Refined grains (e.g., bread, pasta, crackers, baked goods, cereals)	Low-fat dairy (e.g., yogurt, kefir, milk)
Sweets (e.g., desserts, candy, soda)	Vegetables
	Whole grains (e.g., rice, oats, buckwheat, barley)



## Mindfulness

Mindfulness-based practices may help prevent the relapse of depressive episodes. [Mindfulness](#) interventions involve practicing awareness of the present moment. Formal mindfulness practices include body scanning, sitting meditation, and yoga. Informal mindfulness involves integrating mindfulness into daily activities, such as commuting. Consider using a meditation or mindfulness app to help introduce mindfulness practice into your routine.

## Physical activity

Regular aerobic [exercise](#) at moderate or high intensity has been shown to have antidepressant effects. Greater improvements are seen with physical activity led by trained professionals, such as personal training, trainer-led running groups, spin classes, circuit training, and aerobic classes.

## Social support

To reduce feelings of isolation, spend time with and seek support from a trusted friend or family member. Accept help when it is offered by a loved one, and seek out professional guidance in the form of therapy or counseling.



## References

1. Creswell, J. D. (2017). Mindfulness interventions. Annual Review of Psychology, 68, 491–516.
2. Li, Y., Lv, M.-R., Wei, Y.-J., Sun, L., Zhang, J.-X., Zhang, H.-G., & Li, B. (2017). Dietary patterns and depression risk: A meta-analysis. Psychiatry Research, 253, 373–382.
3. National Center for Complementary and Integrative Medicine. (2020, February). Depression. National Institutes of Health. <https://www.nccih.nih.gov/health/depression>
4. National Institute of Mental Health. (2018, February). Depression. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/depression/index.shtml>
5. Public Health Agency of Canada. (2016, December 30). What is depression? Government of Canada. <https://www.canada.ca/en/public-health/services/chronic-diseases/mental-illness/what-depression.html>
6. Qato, D. M., Ozenberger, K., & Olfson, M. (2018). Prevalence of prescription medications with depression as a potential adverse effect among adults in the United States. JAMA: The Journal of the American Medical Association, 319(22), 2289–2298.
7. Quirk, S. E., Williams, L. J., O'Neil, A., Pasco, J. A., Jacka, F. N., Housden, S., Berk, M., & Brennan, S. L. (2013). The association between diet quality, dietary patterns and depression in adults: A systematic review. BMC Psychiatry, 13, 175.
8. Schuch, F. B., Vancampfort, D., Richards, J., Rosenbaum, S., Ward, P. B., & Stubbs, B. (2016). Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. Journal of Psychiatric Research, 77, 42–51.
9. Speed, M. S., Jefsen, O. H., Børglum, A. D., Speed, D., & Østergaard, S. D. (2019). Investigating the association between body fat and depression via Mendelian randomization. Translational Psychiatry, 9(1), 184.



For more educational content and resources: [www.fullscript.com/learn](https://www.fullscript.com/learn)



This handout was developed and medically reviewed by Fullscript's Integrative Medical Advisory team.  
\*These statements have not been evaluated by the Food and Drug Administration. This information is not intended to diagnose, treat, cure, or prevent any disease.

Updated: September 2021